

Multivariable regression on the improvement between baseline and last assessment

ITT-population

Explanator	Regression Coefficient	Standard Error	p
------------	------------------------	----------------	---

Regression on improvement of HAQ			
Gender (female vs male)	-2.62	1.26	0.04
Age (years)	-0.037	0.051	0.47
BMI	0.041	0.105	0.69
Baseline HAQ	0.559	0.067	<0.001
Duration of acute exacerbation > 12 wks	-0.471	1.13	0.68
Initial Dose (5mg vs 10mg)	-0.550	1.09	0.61

Regression on improvement of Three Item Pain Score (TIPS)			
Gender (female vs male)	-1.71	1.67	0.31
Age (years)	-0.003	0.067	0.96
BMI	0.143	0.135	0.29
Baseline TIPS	0.732	0.145	<0.001
Duration of acute exacerbation > 12 wks	-1.48	1.49	0.32
Initial Dose (5mg vs 10mg)	-0.433	1.41	0.76

Multivariable regression on the improvement between baseline and last assessment

ITT-population

Explanator	Regression Coefficient	Standard Error	p
------------	------------------------	----------------	---

Regression on improvement of Arhus Component Pain			
Gender (female vs male)	-2.66	2.82	0.35
Age (years)	-0.017	0.111	0.88
BMI	0.275	0.228	0.23
Baseline Arhus Component Pain	0.613	0.107	<0.001
Duration of acute exacerbation > 12 wks	-2.52	2.50	0.32
Initial Dose (5mg vs 10mg)	0.165	2.35	0.94

Regression on improvement of Arhus Component Invalidity			
Gender (female vs male)	-2.95	1.24	0.02
Age (years)	0.034	0.051	0.50
BMI	0.054	0.104	0.60
Baseline Arhus Component Invalidity	0.402	0.096	<0.001
Duration of acute exacerbation > 12 wks	-1.02	1.11	0.36
Initial Dose (5mg vs 10mg)	-0.297	1.067	0.78