

# A fatal case of aspiration due to consumption of the hallucinogen dipropyltryptamine (DPT)

Merja A. Neukamm, Vanessa Thoma, Susanne Vogt, Volker Auwärter

Institute of Forensic Medicine, Forensic Toxicology, Medical Center – University of Freiburg, Freiburg, Germany



Institute of Forensic Medicine  
Forensic Toxicology

## Background

### Tryptamine-type hallucinogens

- Global drug survey (drug users): 4.2% Tryptamine use
- Serotonin-receptor agonists (5-HT<sub>2A</sub> receptor)
- Non-toxic to organ systems in hallucinogenic doses
- „considerable morbidity“ – due to overreporting?

### Dipropyltryptamine (DPT)

- Adjunct to psychotherapy in 60s / 70s
- Religious sacrament of the “Temple of the True Inner Light”
- Scheduled under the NpSG in 2019
- Typical insufflated dose: 20 to 200 mg



### Effects


#### Tryptamine-type hallucinogens (in general)

- Psychedelic
- Paranoia, psychosis, anxiety,
- Nausea, vomiting
- Hypertension, tachycardia, hyperthermia
- Stimulation (agitation), general tremor

#### Dipropyltryptamine

- Intensity of music and color
- Intense visual and auditory experiences

## Case report



## Analytical results

### Heart blood

- DPT ~ 110 ng/ml
- Sufentanil\*, lidocaine\*, trometamol\*, THC-COOH

### Clinical serum

- DPT 210 ng/ml

### Urine

- DPT ~ 180 ng/ml (detector saturated)
- Lidocaine\*, piperacilline\*



\*Clinical medication

### Stomach

- DPT negative

### All matrices


- Indications of DPT-metabolite Hydroxy-mono-propyltryptamine
- Negative for other drugs, alcohol, synthetic cannabinoids, designer stimulants, other hallucinogens
- Analyses with GC-MS (library search), Toxtyper™, LC-MS/MS



MRM of 4-OH-methyl-ethyl-tryptamine  
but different retention time

## Autopsy

6 days after death



## Autopsy results

- Tongue bites (due to cramps)
- Renal hypoperfusion (shock)
- No needle punctures
- Brain edema (due to oxygen deprivation)

## Discussion

In contrast to most tryptamine overdose reports, there was no agitation, hyperthermia or tachycardia reported in the here presented case. Aspiration of vomit led to blocking of airways, hyperinflation of the lungs and finally to oxygen deprivation of the brain as terminal cause of death. To our knowledge, these are the first published lethal concentrations of dipropyltryptamine.

## Conclusion

Although the young man was reportedly experienced with tryptamine use and although tryptamines usually are relatively non-toxic, death can most likely be attributed to the nasal ingestion of an elevated dose of DPT.

## References

Malaca et al. Int. J. of Mol. Sci. 2020, 21, 9279: Toxicology and Analysis of Psychoactive Tryptamines.

The authors wish to thank Christina Grumann for analysis of DPT in body fluids.

Artwork: commons.wikimedia.org/wiki/File:Psychedelic\_Art\_2022\_The\_Dream\_Entity.png

## Contact

Merja A. Neukamm, Institute of Forensic Medicine,  
Forensic Toxicology, Albertstr. 9, D-79104 Freiburg, Germany  
Fon +49 761 203 6827  
merja.neukamm@uniklinik-freiburg.de